

Mixing Instructions

- Divided into 4-6 portions (Total volume in mls has been pre measured for your pet)
- Mix 1-part dry Critical Care powder with 2 parts warm water. (Double the volume of powder).
Example: 10ml critical care powder with 20ml warm water
- Stir thoroughly to create a well-mixed, lump-free paste. You may adjust the water quantity to reach your desired feeding consistency.

**** You may mix the entire contents of powder with warm water all at once, however, the powder absorbs water over time, so you may need to add a bit more warm water upon each feeding to keep it at the correct consistency.**

Load the syringe

- Remove the plunger of the syringe and spoon 1 portion of the mixture in, then replace the plunger. Ensure you can push the mixture through the syringe. If the mixture is too thick, add more water.

Feeding Instructions

- Slowly give 1 portion by mouth every 4-6 hours over 24-hour period.
- Introduce the tip of the syringe gently into the side of the mouth, in the space between the incisors and molars.
- Push the plunger and administer small amounts from the syringe, pausing after each small amount to allow the pet to swallow.
- Remove the syringe between mouthfuls to prevent choking.
- Continue until the full amount of mixture for that meal has been administered.
- Discard any mixed product after 24 hrs.

How to hold your pet

Wrap your pet in a soft towel so that they are gently restrained and comfortable for the feeding procedure. Have someone assist you if needed.